

Interaction strength, food web topology and the relative importance of species in food webs

Eoin J. O’Gorman^{1,2*}, Ute Jacob^{1,2,3}, Tomas Jonsson⁴ and Mark C. Emmerson^{1,2}

¹Environmental Research Institute, University College Cork, Lee Road, Cork, Ireland; ²Department of Zoology, Ecology and Plant Sciences, University College Cork, Distillery Fields, North Mall, Cork, Ireland; ³Alfred Wegener Institute for Polar and Marine Research, PO Box 120161, D - 27515 Bremerhaven, Germany; and ⁴Ecological Modelling Group, Systems Biology Research Centre, Skövde University, Box 408, SE-54128 Skövde, Sweden

Summary

1. We established complex marine communities, consisting of over 100 species, in large subtidal experimental mesocosms. We measured the strength of direct interactions and the net strength of direct and indirect interactions between the species in those communities, using a combination of theoretical and empirical approaches.

2. Theoretical predictions of interaction strength were derived from the interaction coefficient matrix, which was parameterised using allometric predator–prey relationships. Empirical estimates of interaction strength were quantified using the ln-ratio, which measures the change in biomass density of species A in the presence and absence of species B.

3. We observed that highly connected species tend to have weak direct effects and net effects in our experimental food webs, whether we calculate interaction strength theoretically or empirically.

4. We found a significant correlation between our theoretical predictions and empirical estimates of direct effects and net effects. The net effects correlation was much stronger, indicating that our experimental communities were dominated by a mixture of direct and indirect effects.

5. Re-calculation of the theoretical predictions of net effects after randomising predator and prey body masses did not affect the negative relationship with connectance.

6. These results suggest that food web topology, which in this system is constrained by body mass, is overwhelmingly important for the magnitude of direct and indirect interactions and hence species importance in the face of biodiversity declines.

Key-words: community matrix, dynamic index, indirect effects, press perturbation, trophic links

Introduction

It is now generally accepted that biodiversity has a stabilising effect on natural ecosystems. Diversity has been shown experimentally to have a positive effect on the temporal stability of community biomass (Caldeira *et al.* 2005), decomposition (Dang, Chauvet & Gessner 2005) and parasitism (Tylianakis, Tscharntke & Klein 2006). An increasing number of experimental studies also show that diversity increases the resistance (Kahmen, Perner & Buchmann 2005) and resilience (Steiner *et al.* 2006) of certain ecosystem properties to perturbations. Yet, ecological models based on randomly parameterised community matrices suggest that complexity should destabilise food webs (May 1973; Pimm & Lawton 1978). This implies that real food webs persist despite being unstable and far from equilibrium, or alternatively they have structures and properties that confer stability, in spite of complexity.

The patterning of interaction strengths between predators and prey in natural communities has attracted recent attention. Interaction strengths estimate the magnitude of the effect of one species on another (see Laska & Wootton (1998) for a review of experimental and theoretical approaches to measuring interaction strength). Studies on interaction strength typically measure the direct interaction between two species, but in this study we also examine the net effect of one species on another. The net effect is the sum of the direct and indirect effects of one species on another and can also be taken as a measure of interaction strength. Experiments (Paine 1992; Fagan & Hurd 1994) and theory based on empirical data (McCann, Hastings & Huxel 1998; Neutel, Heesterbeek & de Ruiter 2002) have shown that real food webs are typically characterised by few strong interactions embedded in a majority of weak links. This arrangement of interaction strengths promotes community-level stability by generating negative covariances, which dampen the destabilising potential of strong consumer–resource interactions (McCann 2000; O’Gorman & Emmerson 2009). Although

*Correspondence author. E-mail: e.ogorman@mars.ucc.ie

the detrimental effects of keystone species loss are well documented (Paine 1966; Estes & Palmisano 1974), theoretical studies now suggest that the loss of weak interactors can be damaging for natural communities (Christianou & Ebenman 2005), particularly through their stabilising effects at the landscape level (Berlow 1999; Maser, Guichard & McCann 2007).

Recent theoretical investigations of large empirical food web data sets have also demonstrated the importance of weak interactions in maintaining the rich biodiversity of complex natural communities (Otto, Rall & Brose 2007). It has been predicted that highly connected species could have weaker net effects than poorly connected species in some complex natural ecosystems (Montoya, Emmerson & Woodward 2005). As a result of this property, perturbations to species with fewer links should have larger net effects on the rest of the food web than perturbations to species with many links (Montoya *et al.* 2009). This suggests that complex communities with many highly connected species will be more stable to perturbations than simple communities. However, there is still much doubt as to whether or not this pattern occurs in nature, with several studies indicating that perturbations to highly connected species may in fact destabilise communities (Dunne, Williams & Martinez 2002; Coll, Lotze & Romanuk 2008). Consequently, there is a need to explore empirical patterns of interaction strength and their relationship to species connectance. This paper intends to fill the void of such studies.

The biggest stumbling block towards validating predictions regarding the stabilising patterns of interaction strengths in natural systems is the lack of experimental tests of these theoretical predictions. Many theoretical studies use empirical data to inform their predictions, but theoretical interaction strengths are typically estimated in very different ways to those measured in field experiments (Laska & Wootton 1998; Wootton & Emmerson 2005). This introduces uncertainty over the feasibility of theoretical predictions and constrains their application in the real world. The lack of a direct comparison between theoretical predictions and empirical estimates of interaction strength is a rate limiting step in our understanding of the consequences of biodiversity loss using dynamic systems models.

Testing the validity of theory is essential to confirm that the patterns predicted by community models are indeed the same as those found in nature. Our aim here was to make theoretical predictions about the strength of predator–prey interactions (PPI) in a complex marine community (based on allometric predator–prey relationships) and to compare those predictions to the closest empirical measure of interaction strength that could be determined in a field setting. We chose to use large subtidal exclusion cages to achieve this goal. Exclusion cages have the potential to introduce some experimental artefacts (Benedetti-Cecchi & Cinelli 1997), but they also represent a balance between open and closed systems. Here, they facilitated an intricate exploration of species interactions in a relatively controlled environment, whilst allowing benthic community assembly and ecosystem pro-

cesses to occur against a semi-natural backdrop. We investigated the correlation between our theoretical predictions and empirical estimates of interaction strength (for direct effects and net effects) and searched for common patterns that might contribute to the stability of this complex system.

Materials and methods

STUDY SYSTEM AND EXPERIMENTAL DESIGN

To measure interaction strengths, we established an *in situ* mesocosm experiment in the shallow subtidal of Lough Hyne, a sheltered marine lake in Co. Cork, southwest Ireland (N 51°29'52', W 9°17'46'). Each mesocosm consisted of a subtidal cage, cylindrical in shape, 0.5 m tall, with a diameter of 0.76 m and a 5 mm mesh size (benthic surface area = 0.45 m²). The mesocosms were sealed at both ends and weighted to the benthos by clean, stony substrate, which was spread across the bottom of each cage. This was sufficient to keep the mesocosms in place for the duration of the experiment due to the highly sheltered nature of Lough Hyne. This also meant that the starting conditions for each mesocosm were the same, except for the focal species we selected for manipulation in the experiment. We chose ten abundant benthic species (trophic roles including vertebrate and invertebrate predators, scavengers and grazers) from the shallow subtidal of the Lough, for manipulation in the experiment. The species used were fish [*Gobius niger* (Linnaeus), *Gobius paganelus* (Linnaeus), *Taurulus bubalis* (Euphrasen), *Gaidropsarus mediterraneus* (Linnaeus) and *Ctenolabrus rupestris* (Linnaeus)], decapods [*Carcinus maenas* (Linnaeus), *Necora puber* (Linnaeus) and *Palaemon serratus* (Pennant)] and echinoderms [*Marthasterias glacialis* (Linnaeus) and *Paracentrotus lividus* (Lamarck)]. All of these species are locally common in Lough Hyne, reaching densities in the shallow subtidal during summer months that closely approximate the densities reached in our mesocosms, i.e. 1 individual per 0.45 m² (Costello 1992; Crook, Long & Barnes 2000; Verling *et al.* 2003; Yvon-Durocher *et al.* 2008). The experiment consisted of eleven treatments: ten treatments contained the ten manipulated consumers, with one individual of one species in each cage; the eleventh treatment was a control cage, free of any manipulated species. Note that these were the initial conditions. Once the treatments had been established, the cages were sealed and placed in the shallow subtidal, where benthic invertebrate species small enough to fit through the 5 mm mesh were free to recruit naturally into the cages. There were three replicates of each treatment. This corresponds to a press experiment, where the densities of perturbed species are altered and maintained at predetermined levels throughout the experiment (see Bender, Case & Gilpin (1984) for an overview of press experiments and their application in community ecology). We repeated the experiment in two separate time blocks: 13th July–17th August 2006 (35 days) and 17th August–26th September 2006 (40 days). We used the same cages in each run of the experiment, scrubbing and washing the cages clean before the second experiment. Our aim was to estimate the mean direct effect and the mean net effect of all species within the mesocosm food webs: (i) theoretically, based on allometric predator–prey relationships; and (ii) empirically, based on changes in species biomass density in the presence/absence of all other species in the web.

We used two different sampling substrates to estimate the densities of benthic invertebrate species in the cages: (i) settlement panels, which consisted of 100 × 100 mm PVC squares, were used to estimate the density of sessile species in the mesocosms (sponges, bryozoans, calcareous polychaetes, bivalves); and (ii) nylon pot scourers (c.

radius = 40 mm; *c*. height = 20 mm) were used to estimate the density of mobile species within the mesocosms (amphipods, isopods, gastropods, polychaetes, etc). The pot scourers are ideal for colonisation by benthic invertebrates and simulate the form and structure of coralline algae prevalent in the study system (O’Gorman, Enright & Emmerson 2008). We measured the length of a linear dimension for every individual benthic invertebrate identified from the sampling substrates ($n = 38\,070$). The body mass of each individual was then calculated from length–weight relationships established during the study (see Supporting Information Table S1).

We also noted that two small gobiid fish species [*Gobiusculus flavescens* (Fabricius) and *Pomatoschistus pictus* (Malm)] were able to pass through the small mesh size of the cages. To estimate their densities, we carried out visual surveys on every cage in the experiment, counting the number of gobies seen in each cage over a 5-min period. We also estimated the mean body size of these two species along the south shoreline of the Lough ($n = 100$ for both species). Combining all these data provided an estimate of mean density and mean body mass for most species in each of our 33 mesocosms, for both the August and September experiments. Measuring the density or body mass of resources such as algae, coarse/fine particulate organic matter, and diatoms was problematic. Therefore, we did not take any measurements for the basal resources. This does not affect the way we estimate interaction strength theoretically (see ‘Theoretical predictions of interaction strength’ below), but it meant we could not calculate interaction strengths for the basal resources empirically.

QUANTIFYING FOOD WEB STRUCTURE

We used a combination of gut content analysis and intensive literature research to establish the food web structure of the benthic cages for the August (Fig. S1) and September (Fig. S2) experiments. These composite food webs are built around all the species identified in the experiments (138 taxa in August; 122 taxa in September; see Supporting Information Table S2 for a complete list of taxa). Gut content analysis was carried out on the larger predators in the cages, i.e. manipulated species and the two small gobies ($n =$ at least 30 guts per species). The remaining links were reviewed from the literature. Data from more than 200 publications (peer reviewed journals and books) were standardised using the approach of Martinez (1991), i.e. a direct feeding link was assigned to any pair of species A and B within the benthic cages, whenever an investigator reported that A is likely to consume B in a typical year. This criterion was maintained throughout construction of the web and, for example, restricted the inclusion of prey links that may have occurred by chance through passive consumption. We calculated the number of trophic links for each species in the webs as the number of direct links to the species of interest, including both predators and prey (see Supporting Information Table S2).

THEORETICAL PREDICTIONS OF INTERACTION STRENGTH

The theoretical measures of interaction strength characterising the mesocosm webs were calculated after Montoya *et al.* (2005), i.e. we used theory based on allometric predator–prey relationships to parameterise a community matrix, A , for each of the food webs resolved in our experiments. The community matrix (or interaction coefficient matrix) is derived from the generalised Lotka–Volterra equation and describes the direct effect, a_{ij} , of one individual of a species on the population of another (unlike the Jacobian matrix, which describes the direct effect of the entire population of a species on the total pop-

ulation of another species). The elements of the inverse community matrix following a sign reversal give the sum of the direct and indirect interactions between each species in the food web, i.e. the net effect (Bender *et al.* 1984; Yodzis 1988; Case 2000).

To theoretically predict the direct interaction between a predator and its prey, $^{Th}a_{ij}$, we used the following power function:

$$^{Th}a_{ij} = -b \left(\frac{M_j^{-0.25}}{s_j} \right)$$

where M_j is the body mass of predator j , s_j is the number of prey species consumed by predator j , the intercept b is a free parameter (changing only the magnitude of the interaction strengths) and is set to 0.01 here, and the exponent of -0.25 is based on allometric scaling relationships to approximate basal metabolic rate per unit biomass (West, Brown & Enquist 1997; Brown & Gillooly 2003) (see Supporting Information Appendix S1 for a derivation of this equation). The superscripted Th denotes that this is a theoretically predicted measure of direct effects. Note that $^{Th}a_{ij}$ is derived from a feeding rate, $F_{ij} = -a_{ij}B_i^*B_j^*$, where B_i^* and B_j^* are equilibrium population biomasses of species i and j (see Neutel *et al.* 2002; Emmerson, Montoya & Woodward 2005), so our resulting $^{Th}a_{ij}$ effects are expressed per unit biomass. To predict the direct effect of each prey on its predator, $^{Th}a_{ji}$, we assumed an ecological efficiency, e , equal to 0.1 reflecting a 10% transfer of energy between trophic levels, hence $^{Th}a_{ji} = e^* ^{Th}a_{ij}$ (Brown & Gillooly 2003). Note that the basal resources in the web have no prey, so their only direct effects in the system are calculated using this ecological efficiency. These values comprised the theoretically predicted elements of the community matrix. We then calculated the inverse of the community matrix, whose elements are typically given the notation $(a_{ij})^{-1}$ (Case 2000). By averaging across the absolute values in the columns of the inverse Community matrix, we obtained the mean absolute net effect per unit biomass of each species in the system, $^{Th}(a_{ij})^{-1}$. We calculated mean absolute net effects for both the August and September data sets. Note that the food webs were slightly different for these 2 months (see Supporting Information Figs S1 and S2) and average species body size also varied. Consequently, the theoretical predictions of interaction strength are not identical for the August and September experiments.

EMPIRICAL ESTIMATES OF INTERACTION STRENGTH

We determined the strength of species interactions empirically from our series of press experiments, comparing species changes in the presence/absence of each predator. Since our theoretical predictions of interaction strength deal with effects per unit biomass, we also quantified the empirical measures of interaction strength in units of biomass, calculated as $B = XM$, where X is the density of a given species (m^{-2}) and M is its body mass (mg). We used the dynamic index (Wootton 1997) to empirically measure the direct effect, $^{Em}a_{ij}$, between the manipulated species and each species in the mesocosm food webs:

$$^{Em}a_{ij} = \frac{\ln(B_i^{+j}/B_i^{-j})}{B_j t}$$

where B_i^{+j} and B_i^{-j} are the biomass density of species i in the presence and absence of a predator j , B_j is the biomass density of the corresponding predator and t is the duration of the experiment in days (similar to Brose, Berlow & Martinez 2005). The superscripted Em denotes that this is an empirically estimated measure of direct effects. Note that the change in biomass density was only considered for species pairs where a direct PPI takes place. Ideally, direct interaction

strengths should be measured as the instantaneous growth rate of the focal species, but this is not possible in an experimental setting. Consequently, we divide through by time to obtain the growth rate of the focal species per unit of target species. These $^{Em}a_{ij}$ terms are directly comparable to the terms of the Community matrix, $^{Th}a_{ij}$ defined above. Similar to the community matrix, the dynamic index is derived from the generalised Lotka–Volterra equation (the discrete time version) (Laska & Wootton 1998). An assumption of the dynamic index is that all target species have the same initial densities. This assumption is met, as all mesocosms had the same starting conditions, i.e. empty, apart from the presence or absence of the focal (i.e. manipulated) species.

Measuring direct interaction strength is difficult. The duration of our experiments makes it unclear whether the response of species will be dominated by direct effects, indirect effects or a mixture of the two. At the outset of the experiment, the response of each species will be dominated by direct effects. Over longer timescales indirect effects can feed back in a complex network of interacting species and hence differentiating between direct and indirect effects becomes problematic (Wootton & Emmerson 2005). In our study system, the two experiments lasted for 35 and 40 days. This is insufficient time for entire generations of most species present to occur, but still long enough for indirect effects to become manifest, i.e. the density of a given species was not just affected directly through predation, but also indirectly through trophic (density-mediated) and non-trophic (trait-mediated) interactions (Werner & Peacor 2003; O’Gorman *et al.* 2008). This is illustrated by the presence of positive as well as negative effects of predators on their prey when we calculate $^{Em}a_{ij}$. We also noted changes in the biomass density of species pairs that do not have a direct predator–prey link.

Given the highly reticulate nature of the mesocosm food webs (see Supporting Information Figs S1 and S2), a perturbation to just one species will have knock-on effects for virtually every species in the web. Consequently, we alter our use of the dynamic index to estimate the mean net effect (considering direct and indirect interactions) of each species in the community. First, we consider all species pairs, even if a direct PPI does not take place. Also, we no longer divide through by time. This estimates the long-term change in abundance and is not a growth rate, thus it provides a measure of the net effect of each species, $^{Em}(a_{ij})^{-1}$, combining direct and indirect effects (see Laska & Wootton 1998). As such, these $^{Em}(a_{ij})^{-1}$ effects are comparable to the terms of the negative inverse of the community matrix, $^{Th}(a_{ij})^{-1}$ (Bender *et al.* 1984; Yodzis 1988; Schmitz 1997). It is important to note that in the empirical estimates of interaction strength, species j can have an indirect effect on species i , even if they are not linked along a food chain, because we are accounting for non-trophic interactions (e.g. interference competition, behavioural reactions). This is subtly different from the inverse community matrix, which only takes account of trophic interactions (i.e. direct interactions and density-mediated indirect interactions). As such, we can expect to see some differences in our comparison of theoretical predictions and empirical estimates of interaction strength. If our experiment is dominated by direct effects, the correlation between our theoretical predictions and empirical estimates of direct effects ($^{Th}a_{ij}$ and $^{Em}a_{ij}$, respectively) should be strongest. If our experiment is dominated by net effects, the correlation between our theoretical predictions and empirical estimates of net effects [$^{Th}(a_{ij})^{-1}$ and $^{Em}(a_{ij})^{-1}$, respectively] should be strongest.

The design of the experiment facilitated a comparison of the biomass density of benthic invertebrate species in the presence and absence of the ten manipulated species (by comparing each of the cages containing manipulated species to the mean of the control

cages). However, it was also possible to obtain estimates of the direct/net effect of each benthic invertebrate species on every other species in the mesocosms. Since most benthic invertebrate species were absent from at least some of the cages, we were able to average across the biomass densities of every species in cages where a given benthic invertebrate species was present (B_i^{+}) and compare this to the average of every species in cages where the same benthic invertebrate species was absent (B_i^{-}). In this way, we were able to quantify the effects in each mesocosm attributable to a particular species and thus obtain estimates of the mean absolute direct/net effect of almost all species in the mesocosm webs (some species were present in all treatments and so we could not obtain a comparison with their absence).

EXPLORATION OF BODY MASS PATTERNS

To investigate the relative importance of interaction strength (determined by body mass) vs. food web topology, we randomised the empirical pattern of body masses within the food webs ($n = 1000$ permutations). We calculated per unit biomass effects, using allometric predator–prey relationships to parameterise the community matrix A of theoretically predicted direct effects (as above). We then calculated the predicted mean absolute net effect of each species (averaging across the 1000 permutations) in the inverse community matrix A^{-1} . We also investigated the relationship between body size and number of trophic links, given that we know these food webs to be highly size structured, i.e. big species typically eat smaller species and so they are more likely to be highly connected.

Results

THEORETICAL PREDICTIONS OF INTERACTION STRENGTH

To test for patterns in the theoretically predicted net effects data, we first plotted the frequency distribution of absolute net effects. This reveals a clear skew towards weak effects, with relatively few strong interactions for both the August and September data sets (see Fig. 1a,b). The frequency distribution of direct effects is similarly skewed towards weak effects, but this pattern is not shown. We also plotted the theoretical predictions of the mean absolute direct effect and net effect for each species (a measure of species impact in the system) as a function of the number of trophic links for that species. As number of trophic links increased, the mean direct effect ($^{Th}a_{ij}$) of each species in the web decreased, and this relationship was highly significant for both August (Exponential regression: $F_{1,136} = 467.32$, $P < 0.001$, Fig. 2a) and September (Exponential regression: $F_{1,120} = 419.76$, $P < 0.001$, Fig. 2b). It should be noted that while the relationships in Fig. 2a,b seem to be largely driven by some of the manipulated species (cloud of points in the lower right hand corner of the figure), significant negative relationships remain even if they are removed from the analysis for both August (Exponential regression: $F_{1,126} = 267.60$, $P < 0.001$) and September (Exponential regression: $F_{1,110} = 275.09$, $P < 0.001$). As number of trophic links increased, the mean net effect [$^{Th}(a_{ij})^{-1}$] of each species in the web also decreased, and this relationship was highly significant for




Fig. 1. Frequency distribution of absolute net effects, measured (a, b) theoretically and (c, d) empirically for the two experimental time periods (i.e. the experiment ending in August and the experiment ending in September).




Fig. 2. Mean absolute direct effect (+ SE) of each species on all other species in the mesocosm food webs plotted against number of trophic links (considering both predator and prey links for each species). Direct effects were calculated (a, b) theoretically, using allometric predator–prey relationships and (c, d) experimentally, as growth rates of predators for the two experimental time periods. Values are plotted on a log-log scale with an exponential line of best fit.

both August (Exponential regression: $F_{1,136} = 170.15$, $P < 0.001$, Fig. 3a) and September (Exponential regression: $F_{1,120} = 175.64$, $P < 0.001$, Fig. 3b). Again, highly significant negative relationships remain after the manipulated species are removed from the analysis for both August (Exponential regression: $F_{1,126} = 41.74$, $P < 0.001$) and September (Exponential regression: $F_{1,110} = 55.59$, $P < 0.001$).

EMPIRICAL ESTIMATES OF INTERACTION STRENGTH

We also plotted the frequency distribution of empirically measured net effects. Similar to the theoretical predictions, these are clearly skewed towards weak net effects for both the August and September experiments (see Fig. 1c,d). Again, the frequency distribution of direct effects is similarly skewed towards weak effects, but this pattern is not shown. The




Fig. 3. Mean absolute net effect (\pm SE) of each species on all other species in the mesocosm food webs plotted against number of trophic links (considering both predator and prey links for each species). Net effects were calculated (a, b) theoretically, using allometric predator–prey relationships and (c, d) experimentally, in the presence/absence of predators/competitors for the two experimental time periods. Values are plotted on a log-log scale with an exponential line of best fit.

empirical estimates of mean absolute direct effect ($^{Em}a_{ij}$) were considered as a function of the number of trophic links for each species. There was a significant negative relationship for both the August (Exponential regression: $F_{1,87} = 26.18$, $P < 0.001$, Fig. 2c) and September (Exponential regression: $F_{1,79} = 26.20$, $P < 0.001$, Fig. 2d) experiments. Note that the residual degrees of freedom vary in the empirical relationships because direct effects could not be calculated for all species (see Methods). If we remove the manipulated species from the analysis, the August relationship breaks down (Exponential regression: $F_{1,77} = 2.48$, $P = 0.119$), while a significant negative relationship remains for September (Exponential regression: $F_{1,69} = 6.31$, $P = 0.014$). The empirical estimates of mean absolute net effect [$^{Em}(a_{ij})^{-1}$] were also considered as a function of the number of trophic links for each species. There was a highly significant negative relationship for both the August (Exponential regression: $F_{1,119} = 96.74$, $P < 0.001$, Fig. 3c) and September (Exponential regression: $F_{1,98} = 79.43$, $P < 0.001$, Fig. 3d) experiments. Here, highly significant negative relationships remain even if the manipulated species are removed from the analysis for both August (Exponential regression: $F_{1,109} = 41.89$, $P < 0.001$) and September (Exponential regression: $F_{1,88} = 41.26$, $P < 0.001$).

EXPLORATION OF BODY MASS PATTERNS AND CORRELATIONS

We found a significant correlation between average species body size and number of trophic links for both the August (Pearson's $r = 0.550$, $P < 0.001$) and September (Pearson's

$r = 0.573$, $P < 0.001$) food webs. We also found a significant correlation between the theoretical predictions and empirical estimates of mean absolute direct effects for both August (Pearson's $r = 0.622$, $P < 0.001$; Fig. 4a) and September (Pearson's $r = 0.669$, $P < 0.001$; Fig. 4b). There was a significant, but weaker correlation between total absolute direct effects for both August (Pearson's $r = 0.328$, $P = 0.002$) and September (Pearson's $r = 0.400$, $P < 0.001$). We found an even stronger correlation between the theoretical predictions and empirical estimates of mean absolute net effects for both August (Pearson's $r = 0.877$, $P < 0.001$; Fig. 4c) and September (Pearson's $r = 0.886$, $P < 0.001$; Fig. 4d). Again, there was a significant, but weaker correlation between total absolute net effects for both August (Pearson's $r = 0.661$, $P < 0.001$) and September (Pearson's $r = 0.735$, $P < 0.001$). Lastly, when we compared the interaction strengths from our two experiments, we found significant correlations between the August and September theoretical predictions of $^{Th}a_{ij}$ (Pearson's $r = 0.984$, $P < 0.001$) and $^{Th}(a_{ij})^{-1}$ (Pearson's $r = 0.962$, $P < 0.001$), as well as for the August and September empirical estimates of $^{Em}a_{ij}$ (Pearson's $r = 0.760$, $P < 0.001$) and $^{Em}(a_{ij})^{-1}$ (Pearson's $r = 0.906$, $P < 0.001$) (see Supporting Information Fig. S3 for these correlations). Note that all data were \log_{10} transformed to meet the assumptions of normality for these correlations.

After randomising the pattern of body masses within each food web ($n = 1000$) and re-calculating predicted net effects, we found a significant negative relationship between the theoretically predicted mean absolute net effect of each species and the number of trophic links for both the August




Fig. 4. Correlation between theoretical predictions and empirical estimates of (a, b) mean absolute direct effects and (c, d) mean absolute net effects for the August and September data sets. Values are plotted on a log-log scale with a power fitting.




Fig. 5. Mean absolute net effect (+SE) of each species on all other species in the mesocosm food webs plotted against number of trophic links (considering both predator and prey links for each species). Net effects were calculated theoretically after random rewiring of predator and prey body masses for the two experimental time periods. Values are plotted on a log-log scale with an exponential line of best fit.

(Exponential regression: $F_{1,136} = 85.75$, $P < 0.001$, Fig. 5a) and September (Exponential regression: $F_{1,120} = 100.74$, $P < 0.001$, Fig. 5b) food webs.

Discussion

Our study confirms many of the interaction strength patterns found in other systems. Our theoretical predictions and empirical estimates of net effects show a remarkably similar distribution of interaction strengths, which is highly skewed towards weak effects (see Fig. 1). This arrangement of interaction strengths appears to be a fundamental blueprint for the stability of complex communities (McCann *et al.* 1998; Neutel *et al.* 2002), dampening the destabilising potential of strong consumer-resource interactions (McCann 2000). We also see a clear pattern of lower mean interaction strength (direct effects and net effects) for the more highly connected species, using both empirical estimates and theoretical pre-

dictions (see Figs 2 and 3). Such effects manifest in theoretical explorations of other complex food webs, such as the Ythan Estuary and Broadstone Stream (Montoya *et al.* 2005), but have never been demonstrated empirically. This pattern appears to ensure that complex communities with many highly connected species will be more resilient to perturbations than simple communities with fewer links (Montoya *et al.* 2009), most likely due to an increasing number of pathways for species to dampen impacts of population fluctuations (MacArthur 1955). Dunne *et al.* (2002) and Coll *et al.* (2008) have demonstrated that perturbations to highly connected species have the largest effect on the robustness of model food webs. Although this initially appears to contradict our findings, it should be noted that the perturbation in these studies was extinction of highly connected species. If highly connected species tend to have weaker effects (as we have shown), then removal of these species should increase the mean interaction strength of the community, limiting the

coexistence of many species (Kokkoris *et al.* 2002) and making the food web more sensitive to species removal (Dunne *et al.* 2002).

The pattern appears to be largely driven by the highly size-structured nature of the communities and the fact that body size is correlated to number of trophic links. Here, highly connected species tend to be large and so have small direct/net effects per unit biomass in the community. As such, we argue that the general pattern demonstrated within the Lough Hyne system will be applicable to a wide range of size-structured communities, particularly marine and freshwater systems (Jennings *et al.* 2001; Jonsson, Cohen & Carpenter 2005). We caution that the relationship is less likely to be found in poorly size-structured communities, such as terrestrial food webs, where the direct/net effect per unit biomass of each species is likely to be consistent, irrespective of its connectedness. We demonstrate however, that the relationship between number of trophic links and interaction strength remains even after removing the ten largest species from the current analysis. Here, we obtain two clouds of data in all the panels in Figs 2 and 3 as we are missing information on the direct/net effect of organisms with body sizes intermediate to the large manipulated species and the small benthic invertebrates that assembled through the small mesh size of the cages. Two exceptions are the two small gobies, which do bridge the gap in our regression lines in Figs 2 and 3. With more information on intermediate-sized species, it is likely that the relationships (and r^2 values) would be much stronger.

The weakest patterns are shown for the empirically estimated direct effects. Less than 25% of the variability in number of trophic links is explained by our empirically estimated direct effects. The relationship is weakest for the experiment ending in August, where the negative relationship breaks down with the removal of the ten largest species from the analysis. This suggests that other factors are diluting our estimation of the direct interaction between a predator and its prey, accentuated by the tightly fitting relationship between theoretically predicted direct effects and number of trophic links (see Fig. 2). This is further illustrated by the considerably lower correlation between our theoretical predictions and empirical estimates of direct effects, compared to net effects (see Fig. 4). These results indicate that the duration of our experiments was indeed sufficient for indirect effects to become manifest and that the experimental food webs are dominated by a mixture of direct and indirect effects. Consequently, our empirical measures of net effects offer a clearer insight into the interactions taking place in the system.

It is recognised that biodiversity-ecosystem functioning (BEF) research needs to evolve from a largely confirmatory science to a more predictive discipline (Naeem & Wright 2003). Central to that dictum is the need for unifying patterns in natural systems that can be identified, measured and predicted within the logistical constraints of the natural world. Such an approach will require the integration of sub-disciplines (Ives, Cardinale & Snyder 2005), including BEF research and investigations of PPI, which ultimately describe

food webs. A concerted research effort has demonstrated that the patterning of interaction strengths between predators and their prey is a vital ingredient in the successful functioning of complex ecosystems (McCann *et al.* 1998; Neutel *et al.* 2002; Emmerson & Raffaelli 2004; Montoya *et al.* 2005; O'Gorman & Emmerson 2009). Like so much of BEF research however, there is a large disparity between these studies in the methods used for calculating interaction strength, both within and between theoretical and empirical investigations. To promote an integration of BEF and PPI research as a predictive science, we need to concentrate our efforts on standardising methodologies in all aspects of the discipline.

In this study, we show the predictive power of the community matrix for estimating the strength of species interactions in real food webs. With information on the body size and diet of all the species in our webs, we were able to predict the strength of the interactions (direct and indirect) between all species. The interaction strengths predicted using this modelling approach are correlated to empirical measurements of interaction strength taken in the field over two separate time periods (particularly for the net effects). Importantly, the method we used for predicting interaction strengths theoretically is closely matched to our empirical measurement of interaction strengths (although they do differ in several assumptions, which are discussed below). Schmitz (1997) demonstrated that the inverse community matrix could be used to predict the outcome of field press experiments. Those results, in conjunction with our study, highlight the potential importance of the inverse community matrix in predicting knock-on effects of perturbations in natural food webs, through complex chains of trophic interactions.

Although our study shows the large manipulated species to have the weakest effects per unit biomass in the system, it is also interesting to examine the taxa with the strongest effects. Perhaps surprisingly, predatory polychaetes (particularly the Syllidae and Nereidae), some amphipods (Corophiidae and Dexaminidae), cyclopoid copepods and micro-gastropods had the strongest mean interaction strength per unit biomass. Nereid polychaetes have previously been shown to play an important role in structuring soft-bottom communities (Commito 1982). Deposit feeding amphipods are known to have dramatic effects on the abundance of important basal resources (Gerdol & Hughes 1994). Copepods serve as major grazers of phytoplankton (Turner 2004) and as an important link between microbial and pelagic food webs (Roff *et al.* 1995). Small gastropods can be extremely abundant in shallow marine benthic communities and are likely to have large impacts on the system through deposit feeding and grazing. Little is known about their importance in an overall food web context and their role as strong interactors in the experimental food webs described here highlights the need for detailed study of such species.

It should be noted that there are sources of error involved in the estimation of both the theoretical and empirical effects. For the theoretical predictions, we assumed a 10% transfer of energy between trophic levels to estimate the direct effect

of each prey on its predator. This assumption is based on the second law of thermodynamics, which implies a low efficiency of energy transfer between trophic levels (Brown & Gillooly 2003). Empirical measurements of the transfer efficiency of energy between trophic levels often range from 5% to 15% (Lindeman 1942; Slobodkin 1962). In the absence of more detailed data for individual predator-prey pairs, a 10% transfer of energy is a reasonable approximation. The value of the intercept in the power function is a free parameter. This will not change the qualitative nature of the negative relationship between the net effects and number of trophic links, but it can affect the magnitude of the interaction strengths. Although our theoretical predictions and empirical estimates of direct effects were of the same order of magnitude, our theoretical predictions of net effects tended to be an order of magnitude lower than our empirical estimates of net effects. Changing the value of the intercept would correct this difference. We also accept that the use of a metabolic scaling exponent of -0.25 (derived from $M^{0.75}$ divided through by mass, to express effects in units of biomass; see Supporting Information Appendix S1) can be contentious (Kozłowski & Konarzewski 2004) and works best when spanning body sizes from whales to microbes (West *et al.* 1997). But we contend that the body sizes of organisms in our study still span over 10 orders of magnitude. This is comparable to Tuesday Lake, one of the best defined food webs in the literature, which finds body mass-abundance relationships that agree with a metabolic scaling of 0.75 (or -0.25 per unit biomass) (Jonsson *et al.* 2005).

In the empirical estimates of direct effects, the units of time used in the dynamic index are undefined. We chose to measure the duration of the experiment in days, but the magnitude of the empirical direct effects would be different if we had chosen weeks or hours. Again, this does not alter the qualitative, only the quantitative nature of the results. The empirical estimates of net effects also incorporate non-trophic, as well as trophic, interactions. This is different from the theoretical predictions of net effects, which do not take account of non-trophic interactions in the food webs. It is compelling that we find the same patterns and indeed a strong correlation between the theoretical predictions and empirical estimates of interaction strength, over two separate time periods, in spite of these issues (see Fig. 4). This also suggests that the presence of non-trophic interactions in the empirical measures of interaction strength is not sufficiently important to overwhelm the patterns shown in the theoretical predictions, in spite of previous evidence highlighting the importance of (non-trophic) trait-mediated indirect interactions (Werner & Peacor 2003; Preisser, Bolnick & Benard 2005; O'Gorman *et al.* 2008).

Lastly, it should be noted that net effects measure the combined contribution of every species present in a system on a particular species, in response to a manipulation of another species. Thus, each net effect is uniquely associated with the species composition of the community in which it is measured. We also acknowledge that stochastic processes associated with these complex natural communities will lead to

variation in species composition, which is impossible to control for experimentally. This can lead to misinterpretation of the empirical net effects and probably accounts for much of the unexplained variability in the relationships with the number of trophic links. In addition, our experimental design necessitated that each sub-community began with a different manipulated species in isolation. The extent to which the net effects would change for the aggregate communities shown in Figs S1 and S2 (Supporting Information) is unknown and beyond the scope of the present study. Future studies could examine the extent to which a species' mean net effect in the community changes with the addition or removal of highly connected species.

Interestingly, we find that the pattern of decreasing mean net effect with increasing number of trophic links still holds when we randomise the body sizes used to theoretically predict interaction strength (see Fig. 5). Given two predators of equal size, we would expect the more highly connected predator to have weaker net effects because of s_j in our equation for $^{Th}a_{ij}$. By randomising body size, however, we would expect some cases where highly connected predators are very small and so possess weak net effects. This implies that the strength of direct trophic interactions (and hence the empirical pattern of body mass) is not the factor driving the magnitude of the net effects (Yodzis 1988). Rather, the pattern of net effects seems to be intrinsically linked to the topology of the food web. The definitive test of this would be to randomise topology and see if the pattern between net effects and number of trophic links is destroyed. To randomise topology is to completely change the interacting species and the structure of the food webs under investigation however, and this is no longer a viable comparison with our other estimates of species net effects.

Many studies have quantified the importance of structured hierarchies for the determination of food web topology (Williams & Martinez 2000; Petchey *et al.* 2008). In this context, food web structure reflects the topological arrangement of interaction strengths, which are also known to be size structured. Whilst, the arrangement of direct effects, determined by the empirical pattern of predator and prey body mass, has been shown to affect food web stability (Emmerson & Raffaelli 2004; Brose *et al.* 2005; Otto *et al.* 2007), here the identity of the predators and their prey and their respective body masses also contribute to food web topology. This suggests a deficit in our understanding of food web structure and function and implies that there is a need to integrate dynamic and topological approaches for predicting the behaviour of complex systems, which is currently missing (although see Petchey *et al.* 2008). The approach presented here provides a way of identifying the relative importance of species in a food web context. We have been able to show that there is a strong relationship between empirical and theoretical patterns in the same complex natural food web, which is worthy of further exploration in a wider range of experimental systems. Such a multi-system approach to predicting changes in species importance and the magnitude of species interactions would be an extremely useful tool in conservation biology.

Acknowledgements

We thank Laura Lyons and Órla McLaughlin, for their help in setting up and retrieving the cages used in the experiments. We also thank Owen Petchey and Ulrich Brose for valuable comments on the manuscript. The work was funded by the Irish Research Council for Science Engineering and Technology.

References

- Bender, E.A., Case, T.J. & Gilpin, M.E. (1984) Perturbation experiments in community ecology – theory and practice. *Ecology*, **65**, 1–13.
- Benedetti-Cecchi, L. & Cinelli, F. (1997) Confounding in field experiments: direct and indirect effects of artifacts due to the manipulation of limpets and macroalgae. *Journal of Experimental Marine Biology and Ecology*, **209**, 171–184.
- Berlow, E.L. (1999) Strong effects of weak interactions in ecological communities. *Nature*, **398**, 330–334.
- Brose, U., Berlow, E.L. & Martinez, N.D. (2005) Scaling up keystone effects from simple to complex ecological networks. *Ecology Letters*, **8**, 1317–1325.
- Brown, J.H. & Gillooly, J.F. (2003) Ecological food webs: high-quality data facilitate theoretical unification. *Proceedings of the National Academy of Sciences of the United States of America*, **100**, 1467–1468.
- Caldeira, M.C., Hector, A., Loreau, M. & Pereira, J.S. (2005) Species richness, temporal variability and resistance of biomass production in a Mediterranean grassland. *Oikos*, **110**, 115–123.
- Case, T.J. (2000) *An Illustrated Guide to Theoretical Ecology*, Oxford University Press, New York.
- Christianou, M. & Ebenman, B. (2005) Keystone species and vulnerable species in ecological communities: strong or weak interactors? *Journal of Theoretical Biology*, **235**, 95–103.
- Coll, M., Lotze, H.K. & Romanuk, T.N. (2008) Structural degradation in Mediterranean Sea food webs: testing ecological hypotheses using stochastic and mass-balance modelling. *Ecosystems*, **11**, 939–960.
- Commito, J.A. (1982) Importance of predation by infaunal polychaetes in controlling the structure of a soft-bottom community in Maine, USA. *Marine Biology*, **68**, 77–81.
- Costello, M.J. (1992) Abundance and spatial overlap of gobies (Gobiidae) in Lough-Hyne, Ireland. *Environmental Biology of Fishes*, **33**, 239–248.
- Crook, A.C., Long, M. & Barnes, D.K.A. (2000) Quantifying daily migration in the sea urchin *Paracentrotus lividus*. *Journal of the Marine Biological Association of the United Kingdom*, **80**, 177–178.
- Dang, C.K., Chauvet, E. & Gessner, M.O. (2005) Magnitude and variability of process rates in fungal diversity-litter decomposition relationships. *Ecology Letters*, **8**, 1129–1137.
- Dunne, J.A., Williams, R.J. & Martinez, N.D. (2002) Network structure and biodiversity loss in food webs: robustness increases with connectance. *Ecology Letters*, **5**, 558–567.
- Emmerson, M.C. & Raffaelli, D. (2004) Predator-prey body size, interaction strength and the stability of a real food web. *Journal of Animal Ecology*, **73**, 399–409.
- Emmerson, M.E., Montoya, J.M. & Woodward, G. (2005) Body size, interaction strength and food web dynamics. *Dynamic Food Webs: Multispecies assemblages, ecosystem development, and environmental change* (eds P.C. De Ruiter, V. Wolters & J.C. Moore), pp. 167–178. Academic Press, Amsterdam.
- Estes, A.R. & Palmisano, J.F. (1974) Sea otters – their role in structuring near-shore communities. *Science*, **185**, 1058–1060.
- Fagan, W.F. & Hurd, L.E. (1994) Hatch density variation of a generalist arthropod predator – population consequences and community impact. *Ecology*, **75**, 2022–2032.
- Gerdol, V. & Hughes, R.G. (1994) Effect of *Corophium volutator* on the abundance of benthic diatoms, bacteria and sediment stability in 2 estuaries in southeastern England. *Marine Ecology-Progress Series*, **114**, 109–115.
- Ives, A.R., Cardinale, B.J. & Snyder, W.E. (2005) A synthesis of subdisciplines: predator-prey interactions, and biodiversity and ecosystem functioning. *Ecology Letters*, **8**, 102–116.
- Jennings, S., Pinnegar, J.K., Polunin, N.V.C. & Boon, T.W. (2001) Weak cross-species relationships between body size and trophic level belie powerful size-based trophic structuring in fish communities. *Journal of Animal Ecology*, **70**, 934–944.
- Jonsson, T., Cohen, J.E. & Carpenter, S.R. (2005) Food webs, body size, and species abundance in ecological community description. *Advances in Ecological Research*, **36**, 1–84.
- Kahmen, A., Perner, J. & Buchmann, N. (2005) Diversity-dependent productivity in semi-natural grasslands following climate perturbations. *Functional Ecology*, **19**, 594–601.
- Kokkoris, G.D., Jansen, V.A.A., Loreau, M. & Troumbis, A.Y. (2002) Variability in interaction strength and implications for biodiversity. *Journal of Animal Ecology*, **71**, 362–371.
- Kozłowski, J. & Konarzewski, M. (2004) Is West, Brown and Enquist's model of allometric scaling mathematically correct and biologically relevant? *Functional Ecology*, **18**, 283–289.
- Laska, M.S. & Wootton, J.T. (1998) Theoretical concepts and empirical approaches to measuring interaction strength. *Ecology*, **79**, 461–476.
- Lindeman, R.L. (1942) The trophic-dynamic aspect of ecology. *Ecology*, **23**, 399–417.
- MacArthur, R. (1955) Fluctuations of animal populations, and a measure of community stability. *Ecology*, **36**, 533–536.
- Martinez, N.D. (1991) Artifacts or attributes – effects of resolution on the little-rock lake food web. *Ecological Monographs*, **61**, 367–392.
- Maser, G.L., Guichard, F. & McCann, K.S. (2007) Weak trophic interactions and the balance of enriched metacommunities. *Journal of Theoretical Biology*, **247**, 337–345.
- May, R.M. (1973) *Stability and Complexity in Model Ecosystems*. Princeton University Press, Princeton, NJ.
- McCann, K.S. (2000) The diversity-stability debate. *Nature*, **405**, 228–233.
- McCann, K., Hastings, A. & Huxel, G.R. (1998) Weak trophic interactions and the balance of nature. *Nature*, **395**, 794–798.
- Montoya, J.M., Emmerson, M.C. & Woodward, G. (2005) Perturbations and indirect effects in complex food webs. *Dynamic Food Webs: Multispecies Assemblages, Ecosystem Development, and Environmental Change* (eds P.C. De Ruiter, V. Wolters & J.C. Moore), pp. 369–380. Academic Press, Amsterdam.
- Montoya, J.M., Woodward, G., Emmerson, M.C. & Solé, R.V. (2009) Press perturbations and indirect effects in real food webs. *Ecology*, **90**, 2426–2433.
- Naeem, S. & Wright, J.P. (2003) Disentangling biodiversity effects on ecosystem functioning: deriving solutions to a seemingly insurmountable problem. *Ecology Letters*, **6**, 567–579.
- Neutel, A.M., Heesterbeek, J.A.P. & de Ruiter, P.C. (2002) Stability in real food webs: weak links in long loops. *Science*, **296**, 1120–1123.
- O'Gorman, E.J. & Emmerson, M.C. (2009) Perturbations to trophic interactions and the stability of complex food webs. *Proceedings of the National Academy of Sciences of the United States of America*, **106**, 13393–13398.
- O'Gorman, E.J., Enright, R.A. & Emmerson, M.C. (2008) Predator diversity enhances secondary production and decreases the likelihood of trophic cascades. *Oecologia*, **158**, 557–567.
- Otto, S.B., Rall, B.C. & Brose, U. (2007) Allometric degree distributions facilitate food-web stability. *Nature*, **450**, 1226–1229.
- Paine, R.T. (1966) Food web complexity and species diversity. *American Naturalist*, **100**, 65–75.
- Paine, R.T. (1992) Food-web analysis through field measurement of per-capita interaction strength. *Nature*, **355**, 73–75.
- Petchey, O.L., Beckerman, A.P., Riede, J.O. & Warren, P.H. (2008) Size, foraging, and food web structure. *Proceedings of the National Academy of Sciences of the United States of America*, **105**, 4191–4196.
- Pimm, S.L. & Lawton, J.H. (1978) Feeding on more than one trophic level. *Nature*, **275**, 542–544.
- Preisser, E.L., Bolnick, D.I. & Benard, M.F. (2005) Scared to death? The effects of intimidation and consumption in predator-prey interactions. *Ecology*, **86**, 501–509.
- Roff, J.C., Turner, J.T., Webber, M.K. & Hopcroft, R.R. (1995) Bacterivory by tropical copepod nauplii – extent and possible significance. *Aquatic Microbial Ecology*, **9**, 165–175.
- Schmitz, O.J. (1997) Press perturbations and the predictability of ecological interactions in a food web. *Ecology*, **78**, 55–69.
- Slobodkin, L.B. (1962) Energy in animal ecology. *Advances in Ecological Research*, **1**, 69–101.
- Steiner, C.F., Long, Z.T., Krums, J.A. & Morin, P.J. (2006) Population and community resilience in multitrophic communities. *Ecology*, **87**, 996–1007.
- Turner, J.T. (2004) The importance of small planktonic copepods and their roles in pelagic marine food webs. *Zoological Studies*, **43**, 255–266.
- Tylianakis, J.M., Tscharntke, T. & Klein, A.M. (2006) Diversity, ecosystem function, and stability of parasitoid host interactions across a tropical habitat gradient. *Ecology*, **87**, 3047–3057.
- Verling, E., Crook, A.C., Barnes, D.K.A. & Harrison, S.S.C. (2003) Structural dynamics of a sea-star (*Marthasterias glacialis*) population. *Journal of the Marine Biological Association of the United Kingdom*, **83**, 583–592.

- Werner, E.E. & Peacor, S.D. (2003) A review of trait-mediated indirect interactions in ecological communities. *Ecology*, **84**, 1083–1100.
- West, G.B., Brown, J.H. & Enquist, B.J. (1997) A general model for the origin of allometric scaling laws in biology. *Science*, **276**, 122–126.
- Williams, R.J. & Martinez, N.D. (2000) Simple rules yield complex food webs. *Nature*, **404**, 180–183.
- Wootton, J.T. (1997) Estimates and tests of per capita interaction strength: diet, abundance, and impact of intertidally foraging birds. *Ecological Monographs*, **67**, 45–64.
- Wootton, J.T. & Emmerson, M. (2005) Measurement of interaction strength in nature. *Annual Review of Ecology Evolution and Systematics*, **36**, 419–444.
- Yodzis, P. (1988) The indeterminacy of ecological interactions as perceived through perturbation experiments. *Ecology*, **69**, 508–515.
- Yvon-Durocher, G., Montoya, J.M., Emmerson, M.C. & Woodward, G. (2008) Macroecological patterns and niche structure in a new marine food web. *Central European Journal of Biology*, **3**, 91–103.

Received 18 June 2009; accepted 14 December 2009
Handling Editor: Phil Warren

Supporting Information

Additional Supporting Information may be found in the online version of this article.

Appendix S1. Derivation of the power function used to theoretically predict direct effects.

Table S1. Length–weight relationships used to estimate body mass of all species in the experiments.

Table S2. List of species in the food webs shown in Figs. S1 and S2, with associated trophic heights and number of trophic links.

Fig. S1. Composite mesocosm food web for the experiment ending in August.

Fig. S2. Composite mesocosm food web for the experiment ending in September.

Fig. S3. Correlation between August and September estimates of interaction strength.

As a service to our authors and readers, this journal provides supporting information supplied by the authors. Such materials may be re-organized for online delivery, but are not copy-edited or typeset. Technical support issues arising from supporting information (other than missing files) should be addressed to the authors.